



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The next meeting of the society will be held at Munich in August or September, 1891, and a full attendance by our American colleagues would give them a much better idea of the nature and work of the *Astronomische Gesellschaft* than the above necessarily imperfect sketch, and would serve to bind more closely the ties which unite astronomers on both sides of the ocean.

OBSERVATORY OF KIEL, January 2, 1890.

ON THE ORBIT OF μ^2 HERCULIS (Σ 2220).

BY ARMIN O. LEUSCHNER.

[ABSTRACT.]

The duplicity of μ^1 *Herculis* was discovered by Mr. ALVAN CLARK in the year 1856, with an aperture of $7\frac{3}{4}$ inches. Since then the pair has been regularly observed by the most distinguished observers. In 1879 Dr. W. DOBERCK deduced an orbit of the companion, and arrived at a period of 54.25 years. (*Astr. Nachr.*, 2287.) Recent measures seem to point to a somewhat shorter period. At the suggestion of Mr. S. W. BURNHAM, I have deduced a new orbit, employing therefor the excellent method of Prof. v. GLASENAPP. (*Mon. Not. Royal Astr. Soc.*, March, 1889.) Neglecting a couple of observations made by Messrs. SCHIAPARELLI and BURNHAM in the years 1887 and 1889, I arrived at a period almost identical with that obtained by Dr. DOBERCK. The introduction of the last two observations, however,—the only ones since 1883 accessible to me—renders the representation of the distances by means of the position-angles rather difficult.

The following are Dr. DOBERCK's and my own elements :

ELEMENTS.	DOBERCK.	LEUSCHNER.
Ω	$57^{\circ} 57'$	$62^{\circ} 6'.7$
i	$60^{\circ} 43'$	$67^{\circ} 0'.6$
λ	$156^{\circ} 21'$	$181^{\circ} 59'.0$
e	0.3023	0.2139
a	$1''.46$	$1''.369$
T	1877.13	1880.142
P	54.25	45.39

The observations from which the new orbit has been computed were kindly collected by Mr. BURNHAM. The following residuals were obtained from the interpolating curves:

OBSERVER.	EPOCH.	θ_c	s_c	$\theta_o - \theta_c$	$s_o - s_c$
		^c	["]	^o	["]
Dawes.....	1857.50	63.0	1.75	- 3.7	+ 0.07
Secchi.....	1857.85	64.0	1.75	+ 7.7	- 0.01
Dawes.....	1859.70	67.9	1.69	- 7.5	+ 0.36
O. Struve.....	1860.30	68.7	1.65	- 1.0	- 0.01
O. Struve.....	1862.83	75.1	1.44	+ 3.4	+ 0.06
Dawes.....	1864.43	79.6	1.27	- 2.0	+ 0.54
Engelmann.....	1864.49	80.0	1.27	- 12.5	+ 0.43
Winnecke.....	1864.76	80.3	1.25	- 1.5	+ 0.51
Knott.....	1865.43	83.2	1.17	- 3.6	+ 0.67
Dembowski.....	1865.44	83.4	1.17	- 1.4	+ 0.03
O. Struve.....	1866.68	89.3	1.02	+ 0.2	+ 0.08
O. Struve.....	1868.50	101.7	0.79	- 3.0	+ 0.09
O. Struve.....	1871.52	150.0	0.52	+ 6.8	+ 0.10
O. Struve.....	1873.50	188.8	0.59	- 3.3	+ 0.04
Newcomb.....	1874.49	204.1	0.68	- 0.2	+ 0.07
Gledhill.....	1874.64	0.70	- 0.30
Schiaparelli.....	1875.58	215.6	- 0.4
Newcomb.....	1875.68	216.7	+ 9.2
Hall.....	1875.69	216.7	0.81	+ 3.9	+ 0.37
Holden.....	1875.70	216.7	+ 0.9
Hall.....	1876.59	223.7	0.86	- 0.3	- 0.14
Dembowski.....	1876.68	224.4	0.89	- 8.4	- 0.06
Hall.....	1877.59	230.1	0.98	+ 2.7	- 0.13
Schiaparelli.....	1877.59	230.1	0.98	- 2.2	- 0.18
Dembowski.....	1877.62	230.2	0.98	- 0.3	- 0.10
Burnham.....	1878.48	234.9	1.04	\pm 0.0	+ 0.01
Hall.....	1878.50	234.9	1.05	- 1.1	- 0.17
Burnham.....	1879.45	239.9	1.08	+ 2.8	- 0.18
Hall.....	1879.55	241.0	1.12	- 1.5	- 0.15
Schiaparelli.....	1880.46	245.8	1.07	- 15.6	- 0.37

OBSERVER.	EPOCH.	θ_c	s_c	$\theta_0 - \theta_c$	$s_0 - s_c$
		°	"	°	"
Burnham	1880.47	245.8	1.07	+0.1	-0.11
Hall	1880.70	246.3	1.07	± 0.0	+0.01
Burnham	1881.41	250.1	1.03	+2.0	-0.11
Hall	1881.54	250.6	1.03	-1.5	-0.02
Hall	1882.52	256.4	0.96	+2.7	-0.26
Schiaparelli	1882.60	257.0	+9.8
Burnham	1883.53	263.0	0.86	-0.9	-0.12
Hall	1883.58	263.3	0.85	-1.0	-0.11
Schiaparelli	1883.63	263.7	+11.1
Engelmann	1883.96	0.81	-0.20
Schiaparelli	1887.54	317.9	0.49	± 0.4	± 0.00
Burnham	1889.51	358.4	0.57	-0.5	-0.02

BERKELEY, December, 1889.

ON THE SIMILARITY OF CERTAIN ORBITS IN THE ZONE OF ASTEROIDS.

BY DANIEL KIRKWOOD, LL.D.

The occasional separation of comets into two or more fragments is no longer questioned. It may be asked, however, whether phenomena of this nature have been limited to comets, or whether indications of similar divisions may not be traced in the group of asteroids? This question, suggested some time since,* is one of much interest, but data for its discussion are still far from satisfactory. In order to facilitate the comparison of orbits so as to present at one view the most remarkable resemblances, I take a table of asteroidal elements, arranged in the order of their mean distances, and assume small, admissible limits of difference in the values of the distance, eccentricity, inclination, perihelion and node. In this comparison, however, it is to be borne in mind that the difference in longitude of the perihelion, as also in that of the node,

* *The Asteroids* (p. 48).